Problem 5. a. (10 pts) Evaluate the following indefinite integrals

$$i. \int \cos(3x) dx$$

ii.
$$\int xe^{(x^2)}dx$$

b. (10 pts) Find
$$y(x)$$
 such that $y^1 = \frac{1}{y^3}$ and $y(0) = 1$

Problem 6. (10 pts.) Suppose that $f'(x) = e^{(x^2)}$, and f(0) = 10One can conclude from the mean value theorem that

for which numbers A and B?

Problem 4. (15 pts.) A rocket is launched straight up, and its altitude is $h=10t^2$ feet after t seconds. You are on the ground 1000 feet from the launch site. The line of sight from you to the rocket makes an angle θ with the horizontal. By how many radians per second is θ changing ten seconds after the launch?

h
$$\int_{1000}^{1000} ft$$
 $ton \theta = \int_{1000}^{1000} = \int_{1000}^{1000} ft^2$
 $sec^2 \theta d\theta = \int_{1000}^{1000} ft$

when $t = 10$, $h(\theta) = 10.10^2 = 1000$, ho
 $ton \theta = 1$ so $\theta = \frac{1}{4}$
 $\int_{1000}^{1000} ft d\theta = \int_{1000}^{1000} ft d\theta$
 $\int_{1000}^{1000} ft d\theta = \int_{1000}^{1000} ft d\theta = \int_{10$

Problem 6. (10 pts.) Suppose that $f'(x) = e^{(x^2)}$, and f(0) = 10. One can conclude from the mean value theorem that A < f(1) < B

for which numbers A and B?

$$\frac{f(i)-f(o)}{1-o} = f'(c) \text{ for some } C, \text{ occcl.}$$

$$0 = f(i) = f(o) + f'(c) = 10 + e^{(c^2)}, \text{ occcl.}$$

$$0 = f(i) = f(o) + f'(c) = 10 + e^{(c^2)}, \text{ occcl.}$$

$$0 = f(i) = 10 + e^{(c^2)} < 10 + e^{(c^2)}$$

$$0 = f(i) = 10 + e^{(c^2)} > 10 + 1 = 11$$

$$A = 11$$

$$B = 10 + e$$

Problem 5. a. (10 pts) Evaluate the following indefinite integral
i. $\int \cos(3x) dx = \frac{1}{2} \text{Ain}(5x) + C$

ii.
$$\int ze^{(x^2)}dz = \frac{1}{2} \int e^{u} du = \frac{1}{2} e^{u} + C$$

 $u = \chi^2$
 $du = 2xdx = \frac{1}{2} e^{(\chi^2)} + C$

b. (10 pts) Find y(x) such that $y' = \frac{1}{v^3}$ and y(0) = 1

$$\frac{dy}{dx} = \frac{1}{y^3}$$

$$y^3 dy = 4x$$

$$\frac{1}{4}y^4 = x + C$$

$$\frac{1}{4} = C$$

$$y = (4x+1)^{\frac{1}{4}}$$